VERMONT GEOLOGICAL SURVEY

GEOLOGIC MAP OF VERMONT

PLUTONIC ROCKS

(Including a few dike rocks and volcanic rocks.)

PERMIAN OR TRIASSIC

Upper Permian, or Lower or Middle Triassic — White Mountain plutonic-volcanic series

- Bostonite — *Many small laccoliths and lamprophyre dikes do not show.*
- Biotite and hornblende granites
- Hornblende, biotite, quartz and augite syenites
- Nepheline syenite and pulaskite
- Hornblende-biotite diorite; gabbro
- Essexite
- Volcanic breccia, felsitic tuff, and flows

DEVONIAN

Middle (?) Devonian — New Hampshire plutonic series

- Bethlehem gneiss — *Two-mica granodiorite gneiss.*
- Undifferentiated granitic rocks — *Small dikes and sills do not show.*
- Metadiorite — *Dikes and sills of metagabbro, metadiabase and, meta-andesite, too small to show, are chiefly in the Missisquoi, Athol, and Orfordville formations.*
- Biotite quartz diorite gneiss

ORDOVICIAN

Upper Ordovician — Highgate Springs (?) plutonic series

- Undifferentiated granitic rocks
- Ultramafic rocks — *ub, dunite, peridotite, and serpentinite. us, serpentinite, carbonate rock, talc-carbonate rock and steatite. un, undifferentiated ultramafic rocks.*

PRECAMBRIAN

Upper (?) Precambrian

- Stamford gneiss — *Granitic biotite gneiss with megacrysts of microcline.*
- Undifferentiated gneissic biotite granite, quartz monzonite, and granodiorite

Contact

Local horizon showing structure

Thrust fault — *Location of thrust indicated by structural criteria; sawteeth on upper plate.*

Thrust fault — *Exact location uncertain in places; shown by stratigraphic and paleontologic evidence; sawteeth on upper plate.*

Reverse fault — *Dashed where inferred, bar on upthrown side.*

Normal Fault — *Dashed where inferred, bar and ball on downthrown side.*

QUEBEC - CANADA



EDITING

- W. M. Cady
- C. G. Doll
- J. B. Thompson, Jr.

General: M. P. Billings

0 10 20 30 40 50 miles
SCALE

INDEX MAP OF VERMONT

Shows quadrangle grid, sources of primary geologic information, and (on inset map) areas of regional geologic studies and of editorial responsibility.

PRIMARY GEOLOGIC INFORMATION

1. Hawley, David, 1957, Ordovician shales and submarine slide breccias of northern Champlain Valley in Vermont: Geol. Soc. America Bull., v. 68, p. 55-94; also, references 2, 3, and 14.
2. Erwin, R. B., 1957, The geology of the limestone of Isle La Motte and South Hero Island, Vermont: Vt. Geol. Survey Rept. 11; also, references 3, and 14.
3. Ockerman, Philip and Stauffer, Clinton R., 1950, Cambrian Chazyan series of Champlain Valley, New York and Vermont, and its reefs: Am. Assoc. Petroleum Geologists Bull., v. 43, p. 817-853; also, references 1, 2, and 14.
4. Kay Marshall, 1960, manuscript sketch map; also, 1958, Ordovician-Silurian relations in Champlain Valley, Quebec and Ordovician classification: Am. Jour. Sci., v. 256, p. 65-96, and references 1 and 2.
5. McGerrigle, H. W., 1950, Three geological series in northwestern Vermont: Vt. State Geologist 17th bienn. rept. for 1949-50, p. 191.
6. Shaw, A. B., 1958, Stratigraphy and structure of the St. Albans area, northwestern Vermont: Geol. Soc. America Bull., v. 69, p. 519-567; also, Booth, V. H., 1950, Stratigraphy and structure of the Oak Hill succession in Vermont: Geol. Soc. America Bull., v. 61, p. 1131-1168, and references 6 and 7.
7. Dennis, J. G., in press, The geology of the Enosburg area, Vermont: Vt. Geol. Survey Bull.; also, Booth, V. H., 1950, Stratigraphy and structure of the Oak Hill succession in Vermont: Vt. Geol. Soc. America Bull., v. 61, p. 1131-1168, and references 6 and 7.
8. Doll, C. G., 1944, and Cady, W. M., 1954, portion of Peak quadrangle, manuscript map and notes; C. G. Doll, 1948, manuscript map and notes: Vt.
9. Cady, W. M., Albee, A. L., and Chidester, A. H., in press, Bedrock geology and asbestos deposits of the Lowell Mountain-Belvidere Mountain area, Vermont: U.S. Geol. Survey Bull. 1122-B; also, references 18 and 19; C. G. Doll, 1954, manuscript map and notes: Vt. Geol. Survey.
10. Cady, W. M., 1956, northeast corner of Irasburg quadrangle, Vermont: U.S. Geol. Survey Geol. Quad. Map GQ-74; also, Doll, 1954, manuscript map and notes: Vt. Geol. Survey.
11. Doll, C. G., 1951, Geology of the Memphremagog quadrangle and the southeastern portion of the Irasburg quadrangle, Vermont: Vt. Geol. Survey Bull. 3, 80 p.
12. Goodwin, B. K., in press, Geology of the Island Pond area, Vermont: U.S. Geol. Survey Bull.
13. Cady, W. M., Jr., in press, Geology of the Vermont portion of the Averill quadrangle: Vt. Geol. Survey Bull.; also, Chapman, R. W., 1954, Criteria for the mode of emplacement of alkalic stock at Mount Monadnock, New Hampshire: Geol. Soc. America Bull., v. 65, p. 97-114.
14. Fowler, Phillip, 1950, Geology of the Caledonia quadrangle in Champlain Valley: Am. Jour. Sci., v. 248, p. 1-23; also, references 1, 2, and 3.
15. Stone, S. W., 1953, Structure and stratigraphy of the Milton area, Vermont: manuscript map and text; also, reference 16.
16. Dennis, J. G., 1960, east half of the Milton quadrangle: Vt. Geol. Survey manuscript map and notes; also, Booth, V. H., 1950, Stratigraphy and structure of the Oak Hill succession in Vermont: Geol. Soc. America Bull., v. 61, p. 1131-1168, and reference 6.
17. Christman, R. A., 1961, Geology and of the Mount Mansfield quadrangle, Vermont: Vt. Geol. Survey Bull. 12, 75 p.; also, Booth, V. H., 1950, Stratigraphy and structure of the Oak Hill succession in Vermont: Geol. Soc. America Bull., v. 61, p. 1131-1168.
18. Albee, A. L., 1957, Bedrock geology of the Hyde Park quadrangle, Vermont: U.S. Geol. Survey Geol. Quad. Map GQ-97.
19. Konig, R. H. and Dennis, J. G., in press, The Geology of the Hardwick area, Vermont: Vt. Geol. Survey Bull.
20. Dennis, J. G., 1956, The geology of the Lyndonville area, Vermont: Vt. Geol. Survey Bull. 8, 98 p.
21. Woodland, B. G., in press, The geology of the Burke quadrangle, Vermont: Vt. Geol. Survey Bull.
22. Johansson, W. J., in press, Geology of the Lunenburg quadrangle, Vermont: Vt. Geol. Survey Bull. 14; also,
23. Walley, C. W., 1961, Bedrock geology of the central Champlain Valley of Vermont, Vt. Geol. Survey Bull. 14; also, references 14, 24, and 36.
24. Cady, W. M., 1945, Stratigraphy and structure of westernmost Vermont: Am. Jour. Sci., v. 243-A, p. 497-529, 515-587; also, Osberg, P. H., 1952, The Green Mountain anticlinorium in the vicinity of Rochester and East Middlebury, Vt.: Vt. Geol. Survey Bull. 5, 127 p., and references 23, 37, 38, and 45.
25. Cady, W. M., 1959, eastern areas of the Burlington and Middlebury 15' quadrangle, manuscript maps and notes.
26. Christman, R. A., and Secor, D. T., Jr., 1961, Geology of the Camels Hump quadrangle, Vermont: Vt. Geol. Survey Bull. 13.
27. Cady, W. M., 1956, Bedrock geology of the Montpelier quadrangle, Vermont: U.S. Geol. Survey Geol. Quad. Map GQ-79.
28. Konig, R. H., 1961, Geology of the Plainfield quadrangle, Vermont: Vt. Geol. Survey Bull. 15.
29. Hall, L. M., 1959, The geology of the St. Johnsbury quadrangle, Vermont and New Hampshire: Vt. Geol. Survey Bull. 12, 105 p.
30. Eric, J. H. and Dennis, J. G., 1958, Geology of the Concord-Waterford area, Vermont: Vt. Geol. Survey Bull. 11, 66 p.; also, Billings, M. P., 1937, Regional metamorphism of the Littleton-Moosilauke area, New Hampshire: Geol. Soc. America Bull., v. 48, p. 463-566.
31. Brown, R. L., Jr., 1954, southeast one-ninth of Middlebury quadrangle: manuscript map and text.
32. Cady, W. M., Albee, A. L., and Murphy, J. F., in press, Bedrock geology of the Lincoln Mountain quadrangle, Vermont: U.S. Geol. Survey Geol. Quad. Map GQ-164.
33. Jahns, R. H., and White, W. S., 1960, Bedrock geology and structure of the Chester dome, Vermont: manuscript map and notes; also, references 34 and 35.
34. Murthy, V. R., 1957, Bedrock geology of the East Barre area, Vermont, Vt. Geol. Survey Bull. 10, 121 p.; also, references 33 and 64, White, W. S., and Eric, J. H., 1944, Preliminary report, geology of the Orange County copper district, Vt.: U. S. Geol. Survey open file report.
35. White, W. S., and Billings, M. P., 1951, Geology of the Woodsville quadrangle, Vermont-New Hampshire: Geol. Soc. America Bull., v. 62, p. 647-695; also, reference 64, and White, W. S. and Eric, J. H., 1944, Preliminary report, geology of the Orange County copper district, Vt.: U. S. Geol. Survey open file report.
36. Rosenfeld, J. L., 1960, Barnet quadrangle: manuscript map and notes; also, reference 37.
37. Cady, W. M., and Zen, E-an, 1960, Stratigraphic relationships of the Lower Ordovician Chipman formation in west-central Vermont: Am. Jour. Sci., v. 258, p. 728-739; also, references 14 and 24.
38. Key, Marshall, 1959, Excursions at north end of the Taconic Mountain area, Vermont, in Stratigraphy of structure of west-central Vermont and adjacent New York: 51st New England Intercollegiate Geol. Conf., Rutland, Vt., Guidebook, p. 17-18; also, references 24 and 45.
39. Osberg, P. H., 1959, The stratigraphy and structure of the Core Mountain area, Vermont, in Stratigraphy and structure of west-central Vermont and adjacent New York: 51st New England Intercollegiate Geol. Conf., Rutland, Vt., Guidebook, p. 45-52; also, Fowler, Phillip, 1950, Stratigraphy and structure of the Castleton area, Vermont: Vt. Geol. Survey Bull. 2, 83 p., and references 49 and 48.
40. Osberg, P. H., 1960, Rochester quadrangle and eastern part of Sudbury quadrangle: manuscript map and notes; also, 1952, The Green Mountain anticlinorium in the vicinity of Rochester and East Middlebury, Vt., Vt. Geol. Survey Bull. 5, 127 p., and reference 39.
41. Ern, E. H., Jr., in press, Bedrock geology of the Randolph quadrangle, Vermont: Vt. Geol. Survey Bull.; also, reference 64.
42. Doll, C. G., 1944, A preliminary report on the geology of the Strafford quadrangle, Vermont: Vt. State Geologist 24th bienn. rept. for 1943-1944, p. 14-28; also, White, W. S., and Eric, J. H., 1944, Preliminary report, geology of the Orange County copper district, Vt.: U. S. Geol. Survey open-file report; Howard, P. F., 1959, Structure and rock alteration at the Elizabeth Mine, Vermont: Econ. Geol., v. 54, p. 1214-1249 and 1414-1443, and reference 64.
43. Hadley, J. B., 1950, Geology of the Bradford-Thetford area, Orange County, Vermont: Vt. Geol. Survey Bull. 1, 56 p.; also, 1942, Stratigraphy, structure, and petrology of the Mt. Cube area, New Hampshire-Vermont: Geol. Soc. America Bull., v. 53, p. 113-176; White, W. S., and Eric, J. H., 1944, Preliminary report, geology of the Orange County copper district, Vt.: U. S. Geol. Survey open-file report, and reference 64.
44. Cady, W. M., 1937, 1960, Zen, 1959, portion of Whitehall quadrangle: Vt. Geol. Survey (in part) manuscript maps and notes; also, Rodgers, John, 1937, Stratigraphy and structure in the upper Champlain Valley: Geol. Soc. America Bull., v. 48, p. 1573-1588, and references 23, 24, and 45.
45. Zen, E-an, 1961, Stratigraphy and structure at the north end of the Green Mountain anticlinorium in west-central Vermont: Geol. Soc. America Bull., v. 72, p. 293-338; and, in press, Stratigraphy and structure at a portion of the Castleton quadrangle, Vermont: Vt. Geol. Survey Bull.; also, Theokritoff, George, 1959, Geology of the 'slate belt' in the vicinity of Hampton in Washington County, New York: manuscript map and notes; also, Fisher, D. W., 1961, Stratigraphy and structure of the Taconic rocks in the Hoosick Falls area: Vt. Geol. Survey Bull. 2, 83 p.; Kaiser, E. F., 1945 Northern end of the Taconic mapped sheet in eastern Vermont: Geol. Soc. America Bull., v. 56, p. 1079-1098; and references 24, 36, 37, 38, 46, and 51.
46. Bruce, W. F., 1953, The geology of the Rutland area, Vermont: Vt. Geol. Survey Bull. 4, 136 p.; also, Fowler, Phillip, 1950, Stratigraphy and structure of the Castleton area, Vermont: Vt. Geol. Survey Bull. 2, 83 p., and references 39, 43, and 51.
47. Ern, E. H., Jr., 1960, northern third of the Woodstock quadrangle: Vt. Geol. Survey manuscript map and notes.
48. Chang, P. H., 1950, Structure and metamorphism of the Bridgewater-Woodstock area, Vermont: Manuscript map and text; also, references 46, 47, and 52.
49. Lyons, J. B., 1955, Geology of the Hanover quadrangle, New Hampshire-Vermont: Geol. Soc. America Bull., v. 66, p. 105-148.
50. Shumaker, A. C., 1960, Geology of the Pawlet quadrangle, Vermont: Vt. Geol. Survey manuscript map and text.
51. Thompson, J. B, Jr., 1959, Stratigraphy and structure in the Vermont Valley and the eastern Taconics between Clarendon and Dorset, in Stratigraphy and structure of western Vermont: Vermont Geol. Soc. and 24th New England Intercollegiate Geol. Conf., Rutland, Vt., Guidebook, p. 71-87; also, Fowler, Phillip, 1950, Stratigraphy and structure of the Castleton area, Vermont: Vt. Geol. Survey Bull. 2, 83 p., and references 45, 46, 50, 52, 54, 55, 56.
52. Thompson, J. B, Jr., 1960, Wallingford, Ludlow, and Claremont quadrangles and vicinity, Vt. Geol. Survey (in part): manuscript maps and notes; also, Thompson, J. B., Jr., 1950, A gneiss dome in southeastern Vermont: manuscript map and text; Thompson, J. B., Jr., 1954, Structural geology of the Skitchewaug Mountain area, Claremont quadrangle, Vermont-New Hampshire, 46th New England Intercollegiate Geol. Conf., Dartmouth College, Hanover, p.35-41; Eggleston, J. W., 1918, 1919, Eruptive rocks at Cuttingsville, Vt.: Am. Jour. Sci., 4th ser., v. 45, p. 377-410 and Vt. State Geologist 11th bienn. rept., 1917-1918, p. 167-193, and reference 48.
53. Chapman, R. W. and Chapman, C. A., 1940, Cauldron subsidence at Ascutney Mountain, Vermont: Geol. Soc. America Bull., v. 51, p. 191-211 and Daly, R. A. 1903, Geology of Ascutney Mountain, Vermont: U. S. Geol. Survey Bull. 209, 122 p.
54. Hewitt, P. C., in press, The Geology of the Equinox quadrangle, Vermont and vicinity: Vt. Geol. Survey Bull.
55. Labaluke, R. F., 1960, southeastern third of the Equinox quadrangle: manuscript map and notes (incorporated in Geol. Map of the Manchester quadrangle); Vermont Geol. Survey quadrangle, Vermont and Vicinity: Vt. Geol. Survey Bull.).
56. Wilhelm, D. E., 1959, western part of the Londonderry quadrangle: Vt. Geol. Survey manuscript map and notes; also, Skehan, J. W., reconnaissance notes.
57. Rosenfeld, J. L., 1950, Saxtons River quadrangle and Preliminary report, geology of the Saxtons River-Bellows Falls district, Vt.: U. S. Geol. Survey open-file report, and reference 64.
58. MacFadyen, J. A., Jr., 1956, The geology of the Bennington area, Vermont: Vt. Geol. Survey Bull. 7, 72 p.; also, Prindle, L. M. and Knopf, E. B., 1932, Geology of the Taconic quadrangle: Am. Jour. Sci., 5th ser., v. 24, p. 257-288.
59. Skehan, J. W., in press, The Green Mountain anticlinorium in southern Vermont: Vt. Geol. Survey Bull.; also, Theokritoff, George, 1959, Geology of the 'slate belt' in the vicinity of Hampton in Washington County, New York: manuscript map and notes; also, Fisher, D. W., 1961, Stratigraphy and structure of the Taconic rocks in the Hoosick Falls area: Vt. Geol. Survey Bull. 2, 83 p.; Kaiser, E. F., 1945 Northern end of the Taconic mapped sheet in eastern Vermont: Geol. Soc. America Bull., v. 56, p. 1079-1098; and references 24, 36, 37, 38, 46, and 51.
60. MacDonald, G. J. F., 1960, western two-thirds of the Readsboro quadrangle: manuscript map and notes; also, Rosenfeld, J. L., 1954, Geology of the southern portion of the Chester dome, Vermont: manuscript map and notes; Bulk, Robert, 1956, Bedrock geology of the Massachusetts portion of the Bernardston quadrangle, Massachusetts-Vermont: U. S. Geol. Survey Geol. Quad. Map GQ-90.
61. Segerstrom, Kenneth, 1956, Bedrock geology of the Colrain quadrangle, Massachusetts-Vermont: U. S. Geol. Survey Geol. Quad. Map GQ-86.
62. Moore, G. E., Jr., 1949, Structure and metamorphism of the Keene-Marlboro area, New Hampshire-Vermont: manuscript map and text; also, references 46, 47, and 52.
63. Lyons, J. B., 1955, Geology of the Hanover quadrangle, New Hampshire-Vermont: Geol. Soc. America Bull., v. 66, p. 105-148.
64. Shumaker, A. C., 1960, Geology of the Pawlet quadrangle, Vermont: Vt. Geol. Survey manuscript map and text.
65. Billings, M. P., Rodgers, John and Thompson, J. B. Jr., 1952, Geology of the Appalachian Highlands of east-central New York, southern Vermont, and southern New Hampshire-Field Trip No. 1, in Geol. Soc. America, Guidebook for field trips in New England, p. 1-71.

General: C. G. Doll, 1947-1961, Basic mapping program: Vt. Geol. Survey.

REGIONAL GEOLOGIC STUDIES

63. Cady, W. M., 1960, Stratigraphic and geotectonic relationships in northern Vermont and southern Quebec: Geol. Soc. America Bull., v. 71, p. 531-576.
64. White, W. S., and Jahns, R. H., 1950, Structure of central New England: Jour. Geology, v. 58, p. 179-220.
65. Billings, M. P., Rodgers, John and Thompson, J. B. Jr., 1952, Geology of the Appalachian Highlands of east-central New York, southern Vermont, and southern New Hampshire-Field Trip No. 1, in Geol. Soc. America, Guidebook for field trips in New England, p. 1-71.

General: C. G. Doll, 1947-1961, Basic mapping program: Vt. Geol. Survey.



SECTION ALONG LINE A—A'



SECTION ALONG LINE C—C'



SECTION ALONG LINE E—E'

Williams & Heintz Map Corporation Washington 27, D. C.

EXPLANATION

TACONIC RANGE | CHAMPLAIN AND VERMONT VALLEYS | GREEN MOUNTAINS | EASTERN VERMONT

Middle Ordovician

Trenton group

Pawlet formation (Opw)
Silver gray to jet black, locally carbonaceous and pyritiferous, micaceous silty slate; interbedded, at intervals of a few inches to tens of feet, by beds of dark gray rusty weathered greywacke a few inches to 6 feet thick. The greywacke contains abundant angular grains of quartz, and less abundant feldspar and slate fragments, in a gray argillaceous matrix that is locally calcareous.

Hathaway formation (Oha)
Gray to black argillite and bedded radiolarian chert, with included blocks and fragments of chert, limestone, dolomite, sandstone and greywacke.

Iberville formation (Oi)
Noncalcareous black shale interbedded with occasional dolomite beds and in the lower part with calcareous shale.

Stony Point formation (Osp)
Predominantly calcareous black shale that grades upward into argillaceous limestone and rare dolomite beds, in northeastern Vermont.

Cumberland Head formation (Och)
Interbedded calcareous black shale and fine-grained homogeneous, dark-gray limestone; shown only in Grand Isle county where it is thick enough and well enough exposed to map.

Glens Falls formation (Ogf)
Thin bedded, dark blue-gray, rather coarsely granular and highly fossiliferous limestone.
Ogs. *Shoreham member: interbedded limestone and shale, contains Cryptolithus tessellatus and Prasopora orientalis.*
Ogl. *Larrabee member: thin-bedded shaly limestone.*

Black River group

Orwell limestone, and Isle La Motte and Lowville limestones (Oor, Ol)
Smooth ledged, calcilithographic and lithographic, dove-gray weathered limestone commonly cut by veins of white calcite; the Orwell is a fossiliferous, coarsely clastic limestone. The Lowville is a thin, undifferentiated unit near the base of the Orwell that is characteristically ashen gray and contains abundant Physinopsis tubulosum. The Isle La Motte is about the equivalent of the Orwell in areas west of Champlain thrust, on Isle La Motte and near South Hero, Highgate, Swanton, and St. Albans; it is overlain by the Lowville, which is too thin to show on the map.
Oor. *Rond Pond quartzite member: massive quartz sandstone, near Benson and West Haven, that overlies Orwell limestone.*

UNCONFORMITY

Chazy group

Mount Hamilton formation (Cmh)
White weathered black, gray, green, purple, and red hard slates, some interbedded with thin cherty appearing quartzites and calcareous sandstones; also quartz-pebble and slate bed-conglomerate quartzite, a few inches to several feet thick, locally containing layers of edgewise conglomerate, and a polymict limestone conglomerate. Lithic features vary laterally and are in many places indistinguishable from those of the underlying Hatch Hill and West Castleton formations.

Witnooski dolomite (Cw)
Buff-weathered, pink, buff, and gray dolomite; beds 1 inches to 1 foot thick separated by thin, protruding, red, pink, green, and black siliceous partings.

Monkton quartzite (Cm)
Distinctively red quartzite interbedded with lesser buff and white quartzite and relatively thick sections of dolomite like that of the Witnooski; the quartzites thin to the east, and they become gray and phyllitic to the east and south.

Dunham dolomite (Cd)
Buff-weathered siliceous dolomite, pink and cream mottled or buff to gray on fresh surface; lower part is massive and upper part is sandy and resembles the Witnooski dolomite.

Dalton formation (Cdt)
Schistose quartzite containing pebbles of feldspar and blue quartz; impure dolomite containing pebbles of quartz and feldspar occurs locally. Conglomeratic common near base. Occurs in southwestern Vermont.

Moosalamoo phyllite (Cmo)
Gray to black sericite-quartz phyllite, sericite-quartz-chlorite phyllite occurs locally.

Forestdale marble (Cf)
Buff to rusty-weathered white, buff, and pink and white mottled dolomite containing local interbeds of dolomitic sandstone, gray-green phyllitic quartzite, and crossbedded sandy dolomite.

Bridgeman Hill formation (Cbh)
Undifferentiated dolomite, slate, and conglomerate, on east limb of St. Albans synclinorium, about equivalent to Dunham, Parker, Rugg Brook, and Saxe Brook formations.

Cheshire quartzite (Cc)
Very massive, white to faintly pink or buff siliceous quartzite near the top in eastern and southeastern Vermont; predominantly a vitreous, coarse-sparring mottled gray, somewhat phyllitic quartzite; dolomitic sandstone and conglomerate near the base of the formation in west-central Vermont apparently grades southward into the Dalton formation.

Beekmantown group

Mount Holly complex (pCmh)
Mainly fine- to medium-grained biotitic gneiss, locally migmatitic, and in western areas chloritic; massive and granitoid in some localities, fine-grained or schistose and compositionally layered in others; also abundant amphibolite and hornblendic gneiss, and minor beds of mica schist, quartzite and calc-silicate granulite; includes numerous small bodies of pegmatite and pervasively granitic rock.
pCmq. *quartzite, locally in massive beds as much as 30 feet thick, micaceous quartzite, and quartz-mica schist that commonly contains garnet or pseudomorphs (largely chlorite) after garnet; schists are locally rusty weathered and contain conspicuous flakes of graphite; also includes amphibolite and minor hornblende gneiss, biotite gneiss, and pegmatite.*
pCm. *calcite and dolomite marbles, locally coarse grained; commonly contain phlogopite, actinolite, and diopside, and are interbedded with medium- to coarse-grained calc-silicate granulite; includes minor amounts of other types of Precambrian rock.*

PROFOUND UNCONFORMITY

Gneiss, quartzite, calc-silicate granulite (pCgq)

Upper Cambrian

Hatch Hill and West Castleton formations, undifferentiated (Cch)
The Hatch Hill, a relatively thin formation that succeeds the West Castleton, is characterized by rusty and spongy weathered gray calcareous quartzite traversed by numerous small tubular burrows. The West Castleton is a dark-gray, siliceous, carbonaceous, and pyritiferous slate containing paper-thin white sandy laminae. Black slates are common to both formations. A blue-gray weathered black limestone is near the base of the West Castleton in a few places.

Danby and Potsdam formations (Cdp)
The Danby is comprised of interbedded quartzite and dolomite; white quartzite beds, more than a foot thick, separated by 10 to 12 feet of dolomite in eastern areas, become progressively more dolomitic to the south and west; the massively bedded Potsdam quartzite, west of Orwell thrust.

Saxe Brook dolomite (Csb)
Sandy gray dolomite, sandstone containing dolomitic cement, and pure dolomite; on west limb of St. Albans synclinorium.

Rugg Brook formation (Crb)
Sandy gray dolomite, dolomite conglomerate, and interbeds of gray-weathered sandstone, in St. Albans and Middlebury synclinoria.

UNCONFORMITY

Parker slate (Cpa)
Gray to black micaceous shale and slate, includes dolomite, sandstone, and quartzite lenses; chiefly on west limb of St. Albans synclinorium.

Lower Cambrian

St. Catherine formation (Csc, Csu, Csm, Csc)
Purple, gray-green, and variegated slate and phyllite containing minor interbeds of white to green quartzite; locally schistic. Purple and green chloritoid-bearing slate and phyllite is within dashed line in northern Taconic Range, but not separated farther south.
Csu. *Bomoseen greywacke member: green to olive-colored arkose and greywacke that weathers pale red to white; contains visible flakes of mica and rock fragments.*
Csm. *Zion Hill quartzite member: white weathered green, micaceous chloritic quartzite and greywacke spotted with limonite.*

Breese formation (Cbr)
Dark gray to black phyllite with beds of blue-gray marble, dark gray dolomite, sandy dolomite, and dolomitic sandstone, in upper part; beds of massive quartzite as much as 40 feet thick occur locally and in places contain pebbles of blue quartz. Phyllites are locally highly silistic.
Cbrc. *quartzose green phyllite.*

Precambrian

Granite (Cgr)
Pink, medium-grained, massive, commonly porphyritic; in Barre area.

Granite gneiss (Cgg)
Pink, medium-grained, commonly porphyritic; in Barre area.

Hortonville formation (Oh)
Black, carbonaceous and pyritic slate and phyllite, locally sandy; brown weathering limy beds are common near base. Occurs east of Highgate Springs, Champlain, and Orwell thrusts.

Hortonville, or Cumberland Head, and Glens Falls formations, undifferentiated (Ohg)
Hortonville or Cumberland Head is combined with Glens Falls where the boundary with the Glens Falls is widely covered by surficial deposits, also where the Cumberland Head thins.

Glens Falls and Orwell limestones, undifferentiated (Ogo)
Combined where deformation has made the thin bedded Glens Falls indistinguishable from the thick bedded Orwell; from West Rutland south may contain rocks as low as the Middlebury.

UNCONFORMITY

Middlebury and Chazy limestones; undifferentiated Youngman and Carman formations (Omv, Omd)
Dark blue-gray, somewhat nodular and granular limestone with buff dolomite and shaly interbeds a fraction of an inch thick and 2 to 4 inches apart. The Middlebury, which is east of Champlain and Orwell thrusts, and the Youngman, which is east of Highgate Springs thrust, are, due partly to deformation, more shaly in appearance than the Chazy, which is west of the major thrusts. The Carman is a quartz sandstone with shaly partings that underlies the Youngman. The Chazy contains three members.
Omv. *Valcour member: dark gray calcarenite succeeded by medium- to light gray, buff-weathered silty, partly crystal limestone.*
Omc. *Crown Point member: massive, characterized by abundant Maclurites magnus.*
Omd. *Day Point member: calcareous quartz sandstone, and calcarenite, orange-weathered dolomitic siltstones are common in eastern areas.*

UNCONFORMITY

Chipman, Bridport, and Beldens formations; Providence Island dolomite (Ocbr, Ocb)
The Chipman formation in west-central Vermont includes four members.
Ocbr. *Bridport dolomite member: buff to brown weathered, sharply defined and laterally persistent beds chiefly of medium bedded to massive, scored dolomite, variously designated Bridport formation and Providence Island dolomite in northeastern Vermont.*
Ocbe. *Beldens member: interbedded buff to brown heavily scored dolomite and white to blue-gray marble and limestone; designated Beldens formation east of Highgate Springs thrust.*
Ocw. *Weybridge member: gray limestone with thin interbeds of sandy limestone ½ to 2 inches thick and 1 to 4 inches apart.*
Ocb. *Barnhardts member: blue-gray limestone with irregular spots of light buff dolomite that give weathered surface a mottled appearance.*

Bascom formation, and undifferentiated Luke Hill, Naylor Ledge and Hastings Creek limestones (Obs)
Interbedded dolomite, limestone or marble, calcareous sandstone, quartzite, and limestone breccia; irregular dolomitic layers, thin sandy laminae, and slaty or phyllitic partings characterize limestone and marble of lower, middle, and upper parts of the Bascom, respectively, south of West Rutland it includes some of Chipman formation. The combined Luke Hill, Naylor Ledge, and Hastings Creek, east of Philipsburg thrust, are stratigraphically equivalent to the Bascom
Obh. *Brownell Mountain phyllite member: calcareous phyllite in upper part of the Bascom formation of east limb of Hinesburg synclinorium.*

Cutting dolomite, and undifferentiated Morgan Corner and Wallace Creek formations (Oc)
Typical Cutting is a massive, gray weathered, nonlaminated dolomite with a finely laminated calcareous sandstone at base. The combined Morgan Corner and Wallace Creek formations, east of Philipsburg thrust, are stratigraphically equivalent to the Cutting.

Shelburne, Whitehall, and Stirling Pond formations (Os)
The Shelburne is chiefly a white marble or gray limestone characterized by ruined reticulate lines of gray dolomite on the weathered surface; includes Sutherland Falls marble, intermediate dolomite and Columbian marble of the marble quarries. Interbedded massive dolomite increases westward and predominates in the Whitehall formation, west of Champlain and Orwell thrusts. The Stirling Pond, which is identical to the Shelburne, is east of Philipsburg thrust.

Clarendon Springs, Tironderoga, and Rock River dolomites; Gorge formation (Ocs)
Fairly uniform, massive, smooth weathered gray dolomite characterized by numerous grades and knots of white quartz; quartz sandstone and irregular masses of chert are common in the Clarendon Springs west of Orwell and Champlain thrusts and the Rock River east of Philipsburg thrust. The Gorge is a partly conglomeratic facies on west limb of the St. Albans synclinorium.

Highgate formation (Ohi)
Banded blue limestone and calcareous slate; local lenses of limestone conglomerate, on west limb of St. Albans synclinorium.

Sweetsburg formation (Csh, Cv, Csm, Cs)
Black, carbonaceous, well-cleaved slate with characteristic thin (1-2mm) whitish silty bedding laminae; in St. Albans and Hinesburg synclinoria and Cambridge syncline.
Csh. *Hungerford slate member: black slate.*
Cv. *Rockledge conglomerate member: phenoclasts chiefly of bioformal limestone in a matrix of gray limestone containing frosted quartz sand grains.*
Csm. *Skeels Corners slate and Mill River conglomerate members undifferentiated: black slate; local dolomite, sandstone, dolomite conglomerate, limestone bioherms, and limestone conglomerate lenses. The Mill River is a basal limestone conglomerate.*
Cs. *St. Albans slate member: black, gray-black, or tan micaceous slate.*

Missisquoi formation (Omc, Omcc, Omss, Omq, Omnb, Omm, Omw)
NORTHERN VERMONT
Omcc. *Crom Hill member: pale greenish-gray to black phyllite grades locally into gray to black slate; minor to mafic volcanic rocks.*
Omhb. *Hortons Bridge quartzite member: buff to pale-green quartzite with interbeds of quartz-sericite-chlorite phyllite.*
Omcc. *Coburn Hill volcanic member: actinolite-epidote-chlorite rock, greenstone, and hornblende-albite-epidote amphibolite; also includes slate.*
Omc. *carbonaceous phyllite and slate, black to light gray (where extensive enough to show on map).*
Omq. *Limekiln Hill member: quartz and slate pebble, phyllitic conglomerate with interbeds of slate and phyllite —chiefly quartz-sericite-magnetite-chloritoid rocks.*

SOUTHERN VERMONT
Omc. *Rusty weathering carbonaceous mica schist, quartzite and micaceous quartzite*
Omb. *Barnard volcanic member: fine- to medium-grained biotite gneiss, hornblende gneiss, and amphibolite.*
Omw. *Whetstone Hill member: carbonaceous black to light gray phyllite and schist containing porphyroblasts of biotite and garnet; beds of gray micaceous quartzite, fine-grained biotite gneiss and amphibolite*

NORTHERN AND SOUTHERN VERMONT
Omm. *Monkton member: quartzite and quartz-plagioclase granulite, in layers ½ to several inches thick, separated by "pinstripe" partings that contain muscovite, chlorite, epidote, biotite, and locally garnet; also greenish quartz-sericite-chlorite phyllite and schist; and minor carbonaceous phyllite. Schist and phyllite commonly contain biotite and garnet porphyroblasts in southern Vermont.*

UNCONFORMITY

Underhill formation (Cu)
Silvery, gray-green, quartz-sericite-albite-chlorite-biotite schist containing abundant lenticular segregations of granular white quartz; locally quartz-sericite-albite-chlorite phyllite; porphyroblasts of albite, garnet, and magnetite are common and locally very abundant in gneissic facies in areal anticlines of the Green Mountain anticlinorium.
Cuf. *Fairfield Pond member: greenish quartzitic schist (quartz-sericite-albite-chlorite-biotite); sericite-quartz-chlorite phyllite, locally purple or red, common in lower part.*
Cug. *gneissone: varied composition including albite-chlorite-epidote-calcite and sericite-magnetite-chloritoid-muscovite rocks.*
Cuc. *carbonaceous quartz-sericite-albite-chlorite schist and phyllite.*
Cuw. *White Brook member: chiefly brown-weathered whitish, tan and gray sandy dolomite, locally only a hematitic zone; includes carbonaceous crystalline limestone in Cambridge syncline.*

Pinnacle formation (Cp)
Schistose greywacke, gray to buff, commonly striped, quartz-albite-sericite-biotite-chlorite rock predominates; quartz-cobble and boulder conglomerate is common, chiefly near base.
Cpt. *Tibbit Hill volcanic member: albite-actinolite-chlorite-epidote greenstone; locally pillowed and vesicular.*

Highgate formation (Ohi)
Banded blue limestone and calcareous slate; local lenses of limestone conglomerate, on west limb of St. Albans synclinorium.

Stowe formation (OCsc, OCsg, OCs)
Quartz-sericite-(muscovite-paragonite)-chlorite phyllite and schist; porphyroblasts of albite, garnet, chloritoid, or kyanite are common locally. Includes phyllitic greywacke north of Lamoille River. Schist contains abundant segregations of granular white quartz.
OCsc. *carbonaceous schist and phyllite north of Lamoille River; occurs in several small areas to the south, not shown on map.*
OCsg. *greenstone and amphibolite; epidote-albite-chlorite rocks contain actinolite and hornblende where more metamorphosed.*

Ottauquechee formation (Oq)
Black carbonaceous phyllite or schist containing interbeds of massive quartzite commonly criss-crossed by veins of white quartz; quartzite is dark gray and carbonaceous, light gray, or white; also includes light green quartz-sericite-chlorite phyllite or schist and sericitic quartzite; beds of phyllitic greywacke and feldspar granule conglomerate are common in Granville Basin. Schist contains abundant porphyroblasts of garnet and biotite from Ludlow south.
Cg. *greenstone and amphibolite.*

Pinney Hollow, Ottauquechee, and Stowe formations, undifferentiated (OCq)
Includes quartz-muscovite-garnet-chlorite-biotite schist, rusty carbonaceous schist, amphibolite, and schistose quartzite, schist locally contains prophyroblasts of staurolite and kyanite; on flanks of Chester and Athens domes

Littleton formation (S)
Gray slate and phyllite containing interbeds of gray schistose quartzite ¼ inch to 6 inches thick. West of Guildhall are lustrous, light to dark gray biotite-garnet phyllite and schist, some slate, and subordinate quartzite and impure quartzite. South of Bellows Falls gray phyllite passes eastward into gray mica schist containing porphyroblasts of biotite, garnet, and staurolite.

Fitch formation (Sf)
NORTHEASTERN VERMONT
Bluish-gray, white-weathered, sandy limestone.
SOUTHEASTERN VERMONT
Quartz-plagioclase-biotite granulite; actinolite-diopside granulite; impure limestone and dolomitic mica schist; the carbonate-rich beds are typically an inch or two thick and segmented so as to give the weathered outcrop a characteristic pitted appearance.

Clough formation (Sk)
NORTHEASTERN VERMONT
Boulders, cobbles, pebbles and angular fragments of quartzite, micaceous quartzite, and gray mica schist in matrix of mica schist, quartz-mica schist or quartzite; schist commonly contains porphyroblasts of biotite, less commonly garnet.
SOUTHEASTERN VERMONT
Quartzite, quartz-conglomerate, and mica schist; lenses of fossiliferous calcareous quartzite in upper part.

UNCONFORMITY

Partridge formation (Dp)
NORTHEASTERN VERMONT
Dull gray to rusty black carbonaceous slate and phyllite, mostly fissile, but locally massive and tough. Contains a few beds of both schistose and massive soda-rhyolite tuff (not shown on map).
SOUTHEASTERN VERMONT
Rusty weathering carbonaceous mica schist locally containing porphyroblasts of biotite, garnet, or staurolite.
Dpv. *volcanics: fine-grained biotite gneiss and amphibolite.*

Ammonoosuc volcanics (Oa)
NORTHEASTERN VERMONT
Soda-rhyolite tuff, breccia, and flows.
SOUTHEASTERN VERMONT
Fine-grained chloritic and biotitic gneiss and greenstone in areas north of Bellows Falls, biotite gneiss and amphibolite south of Bellows Falls.

Albee formation (Dal)
Massive, gray, white-weathered quartzite and feldspathic quartzite interbedded with greenish-gray slate, phyllite, feldspathic phyllite and quartzose orgulaceous phyllite. Micaceous quartzite, quartz-mica schist, mica schist, and dark-gray carbonaceous phyllite are common. Garnet, staurolite, and kyanite are in the extremity of granitic plutons. Soda-rhyolite tuff occurs locally. Micaceous quartzite characterized by thin, schistose "pinstripe" partings is common in many areas.

Orfordville formation (Dco, Dof, Dov)
Carbonaceous phyllite; minor quartzite.
Dov. *Sunday Mountain volcanics: greenstone, chloritic schist, felsite, and quartz-feldspar-sericite schist.*
Dop. *Pine Pond volcanics: greenstone and green chloritic schist commonly interbedded with lenses of fine- and coarse-fragmental volcanic schist and conglomerate; chloritic and biotitic gneiss; all west of Ammonoosuc fault; mainly amphibolite east of the Ammonoosuc fault.*

Silurian

Shaw Mountain formation (Ss)
NORTHERN VERMONT
Chiefly tan to brown weathered quartzose limestone and calcareous quartzite characterized by specks of limonite after ankerite; locally underlain by quartz conglomerate and overlain by blue fossiliferous crystalline limestone, greenstone and quartz-sericite schist.
SOUTHERN VERMONT
Quartzite, quartz conglomerate, rummingtonite schist, amphibolite, and quartz-sericite schist with porphyroblasts of biotite and garnet.

UNCONFORMITY

Northfield formation (DSn)
Dark gray to black quartz-sericite slate or phyllite with fairly widely-spaced interbeds a few inches thick of siltstone and silty crystalline limestone like that of the Waits River formation; calcareous slate north of Lamoille River; phyllite passes into gray quartz-sericite schist containing abundant porphyroblasts of biotite and garnet in southern Vermont.

Waits River formation (Dwb, Dwg, Dwm, Dwf, Dwc)
Gray quartzose and micaceous crystalline limestone weathered to distinctive brown earthy crust; interbedded and intergradational with gray quartz-muscovite phyllite or schist. Where more metamorphosed the limestones contain actinolite, hornblende, zoisite, diopside, wollastonite, and garnet, and the phyllite and schist, biotite, garnet, and locally kyanite, andalusite, or staurolite.
Dwc. *Standing Pond volcanic member: amphibolite, garnet amphibolite, coarse garnet schist with fasciculite hornblende, and hornblende massifs; contains pillow lavas near St. Johnsbury and passes eastward into actinolitic greenstone and greenstone south of Windsor.*
Dwf. *Crow Hill member: tough gray quartzite near St. Johnsbury (not separated from other rock that shown on map) occurs at several other places.*
Dwb. *Barton River member: interbedded siliceous crystalline limestone and arkosic-quartzo-chloritic phyllite in northern Vermont; diopsidic limestone and cordierite hornfels at contacts with granitic dikes and sills.*
Dwa. *Ayers Cliff limestone member: siliceous crystalline limestone containing thin beds of slate and phyllite north of the Lamoille River.*

Gile Mountain formation (Dgs, Dgm, Dgh, Dgl)
Gray quartz-muscovite phyllite or schist, interbedded and intergradational with gray micaceous quartzite (greywacke northeast of Vaiehgan River), calcareous mica schist, and, locally, quartzose and micaceous crystalline limestone like that of the Waits River formation. The phyllite and schist commonly contain porphyroblasts of biotite, garnet, or staurolite, and locally kyanite, andalusite, or sillimanite.
Dgs. *Stillwater member: highly feldspathic grit, probably arkosic; feldspathic chloride-andesite schist and amphibolite, all northeast of Nulhegan River.*
Dga. *amphibolite: hornblende-quartz-biotite-chlorite rock.*
Dgm. *Mossalamoose slate member: chiefly gray slate or phyllite characterized by beds of gray schistose quartzite ¼ inch to 3 inches thick.*

Ordovician

Missisquoi formation (Omc, Omcc, Omss, Omq, Omnb, Omm, Omw)
NORTHERN VERMONT
Omcc. *Crom Hill member: pale greenish-gray to black phyllite grades locally into gray to black slate; minor to mafic volcanic rocks.*
Omhb. *Hortons Bridge quartzite member: buff to pale-green quartzite with interbeds of quartz-sericite-chlorite phyllite.*
Omcc. *Coburn Hill volcanic member: actinolite-epidote-chlorite rock, greenstone, and hornblende-albite-epidote amphibolite; also includes slate.*
Omc. *carbonaceous phyllite and slate, black to light gray (where extensive enough to show on map).*
Omq. *Limekiln Hill member: quartz and slate pebble, phyllitic conglomerate with interbeds of slate and phyllite —chiefly quartz-sericite-magnetite-chloritoid rocks.*

SOUTHERN VERMONT
Omc. *Rusty weathering carbonaceous mica schist, quartzite and micaceous quartzite*
Omb. *Barnard volcanic member: fine- to medium-grained biotite gneiss, hornblende gneiss, and amphibolite.*
Omw. *Whetstone Hill member: carbonaceous black to light gray phyllite and schist containing porphyroblasts of biotite and garnet; beds of gray micaceous quartzite, fine-grained biotite gneiss and amphibolite*

NORTHERN AND SOUTHERN VERMONT
Omm. *Monkton member: quartzite and quartz-plagioclase granulite, in layers ½ to several inches thick, separated by "pinstripe" partings that contain muscovite, chlorite, epidote, biotite, and locally garnet; also greenish quartz-sericite-chlorite phyllite and schist; and minor carbonaceous phyllite. Schist and phyllite commonly contain biotite and garnet porphyroblasts in southern Vermont.*

Cavendish formation (OCc, Ccm)
(May in large part correlate with Tyson formation, or may be older, in northernmost area, possibly age equivalent of rocks elsewhere included in Mount Holly complex.)
¹Ccr. *Readsboro member: quartz-muscovite schist containing biotite or chlorite and characterized by conspicuous porphyroblasts of mafic plagioclase; less commonly quartz-muscovite-paragonite schist containing chlorite, garnet, or chloritoid, or, in Chester dome, quartz-muscovite-paragonite schist containing garnet, staurolite, and locally kyanite (Gassetts schist).*
²Ccg. *Bull Hill gneiss: quartz-plagioclase-microline-biotite gneiss characterized in many areas by augen of microline as much as 2 inches long; fine- to medium-grained quartz-plagioclase-biotite or biotite-muscovite gneiss.*
³Ccm. *buff dolomite; minor white to pink calcite marble, actinolitic and diopsidic marbles and beds of actinolite diopside granulite common in Chester dome.*

PROFOUND UNCONFORMITY

Mount Holly complex (pCmh, pCmq, pCm)
Mainly fine- to medium-grained biotitic gneiss, locally migmatitic, and in western areas chloritic; massive and granitoid in some localities, fine-grained or schistose and compositionally layered in others; also abundant amphibolite and hornblendic gneiss, and minor beds of mica schist, quartzite and calc-silicate granulite; includes numerous small bodies of pegmatite and pervasively granitic rock.
pCmq. *quartzite, locally in massive beds as much as 30 feet thick, micaceous quartzite, and quartz-mica schist that commonly contains garnet or pseudomorphs (largely chlorite) after garnet; schists are locally rusty weathered and contain conspicuous flakes of graphite; also includes amphibolite and minor hornblende gneiss, biotite gneiss, and pegmatite*
pCm. *calcite and dolomite marbles, locally coarse grained; commonly contain phlogopite, actinolite, and diopside, and are interbedded with medium- to coarse-grained calc-silicate granulite; includes minor amounts of other types of Precambrian rock.*

Cambrian (?)

Underhill formation (Cu)
Silvery, gray-green, quartz-sericite-albite-chlorite-biotite schist containing abundant lenticular segregations of granular white quartz; locally quartz-sericite-albite-chlorite phyllite; porphyroblasts of albite, garnet, and magnetite are common and locally very abundant in gneissic facies in areal anticlines of the Green Mountain anticlinorium.
Cuf. *Fairfield Pond member: greenish quartzitic schist (quartz-sericite-albite-chlorite-biotite); sericite-quartz-chlorite phyllite, locally purple or red, common in lower part.*
Cug. *gneissone: varied composition including albite-chlorite-epidote-calcite and sericite-magnetite-chloritoid-muscovite rocks.*
Cuc. *carbonaceous quartz-sericite-albite-chlorite schist and phyllite.*
Cuw. *White Brook member: chiefly brown-weathered whitish, tan and gray sandy dolomite, locally only a hematitic zone; includes carbonaceous crystalline limestone in Cambridge syncline.*

Pinnacle formation (Cp)
Schistose greywacke, gray to buff, commonly striped, quartz-albite-sericite-biotite-chlorite rock predominates; quartz-cobble and boulder conglomerate is common, chiefly near base.
Cpt. *Tibbit Hill volcanic member: albite-actinolite-chlorite-epidote greenstone; locally pillowed and vesicular.*

Hazens Notch formation (Ch)
Interbedded carbonaceous and noncarbonaceous quartz-sericite-albite-chlorite schist; grades to quartzite and gneiss.
Chb. *Belvidere Mountain amphibolite member: coarse- to fine-grained hornblende-epidote-albite rock; grades to epidote-chlorite-actinolite-albite greenstone where less metamorphosed.*
Chm. *sericite-quartz-chlorite-albite-magnetite schist, near Hyde Park.*
Chg. *chiefly albite-actinolite-chlorite-epidote greenstone; locally hornblende-epidote-chlorite-albite amphibolite.*

UNCONFORMITY

Cavendish formation (OCc, Ccm)
(May in large part correlate with Tyson formation, or may be older, in northernmost area, possibly age equivalent of rocks elsewhere included in Mount Holly complex.)
¹Ccr. *Readsboro member: quartz-muscovite schist containing biotite or chlorite and characterized by conspicuous porphyroblasts of mafic plagioclase; less commonly quartz-muscovite-paragonite schist containing chlorite, garnet, or chloritoid, or, in Chester dome, quartz-muscovite-paragonite schist containing garnet, staurolite, and locally kyanite (Gassetts schist).*
²Ccg. *Bull Hill gneiss: quartz-plagioclase-microline-biotite gneiss characterized in many areas by augen of microline as much as 2 inches long; fine- to medium-grained quartz-plagioclase-biotite or biotite-muscovite gneiss.*
³Ccm. *buff dolomite; minor white to pink calcite marble, actinolitic and diopsidic marbles and beds of actinolite diopside granulite common in Chester dome.*

PROFOUND UNCONFORMITY

Mount Holly complex (pCmh, pCmq, pCm)
Mainly fine- to medium-grained biotitic gneiss, locally migmatitic, and in western areas chloritic; massive and granitoid in some localities, fine-grained or schistose and compositionally layered in others; also abundant amphibolite and hornblendic gneiss, and minor beds of mica schist, quartzite and calc-silicate granulite; includes numerous small bodies of pegmatite and pervasively granitic rock.
pCmq. *quartzite, locally in massive beds as much as 30 feet thick, micaceous quartzite, and quartz-mica schist that commonly contains garnet or pseudomorphs (largely chlorite) after garnet; schists are locally rusty weathered and contain conspicuous flakes of graphite; also includes amphibolite and minor hornblende gneiss, biotite gneiss, and pegmatite*
pCm. *calcite and dolomite marbles, locally coarse grained; commonly contain phlogopite, actinolite, and diopside, and are interbedded with medium- to coarse-grained calc-silicate granulite; includes minor amounts of other types of Precambrian rock.*

Precambrian

Granite (Cgr)
Pink, medium-grained, massive, commonly porphyritic; in Barre area.

Granite gneiss (Cgg)
Pink, medium-grained, commonly porphyritic; in Barre area.

QUEBEC – CANADA

NEW YORK | NEW HAMPSHIRE

METAMORPHIC ZONES

EXPLANATION

Post Metamorphic Faults

White Mountain Plutonic Series (W)
Typically surrounded by a zone several hundred yards wide of finely andalusite-cordierite or sillimanite-bearing hornfels. These aureoles are distinctly younger than the regional metamorphism.

New Hampshire Plutonic Series (N)
Most bodies emplaced during the regional metamorphism or shortly thereafter. Sillimanite and locally cordierite occur near many contacts in northern part of state in zones too narrow to show on map.

Sillimanite Zone
Marked by first appearance of sillimanite in rocks of appropriate composition.

Staurolite-Andalusite Zone (SA)
Marked by first appearance of staurolite or andalusite in pelitic schists. Some kyanite occurs in southeast part of this zone.

Staurolite-Kyanite Zone (SK)
Marked by first appearance of staurolite or kyanite in pelitic schists. Carbonate rocks commonly contain diopside.

Chloritoid-Kyanite Zone (CK)
Characterized by the assemblage chloritoid-kyanite in northern Green Mountains. Staurolite is absent.

Garnet Zone (G)
Marked by the first appearance of almandite garnet in pelitic schists. Mafic rocks commonly contain a dark green amphibole.

Biotite Zone (B)
Marked by approximate first appearance of biotite in rocks with a low alumina-alkali ratio or of chloritoid in rocks, mainly pelitic phyllites, with a high alumina-alkali ratio. Mafic rocks mainly greenstone schists, though locally containing an actinolitic amphibole. Carbonate rocks may contain talc or phlogopite and locally tremolite or scapolite.

Chlorite Zone (Ch)
Rocks showing evidence of rank somewhat recrystallization but containing few minerals not found in sedimentary rocks. Stilpnomelane is reported locally in schistose greywackes.

Polymetamorphic areas
Mainly rocks metamorphosed in the Taconian orogeny to probable staurolite zone or higher, then later subjected to a low-grade metamorphism (Acadian?). Grade of Paleozoic metamorphism indicated by color tone, ash stippling, and by letters preceding "R" in letter symbol.

Scale of miles 0 10 20 30

METAMORPHIC MAP of VERMONT

QUEBEC – CANADA

FORELAND | GREEN MOUNTAIN ANTICLINORIUM | CONNECTICUT VALLEY – GASPE SYNCLINORIUM

NEW YORK | NEW HAMPSHIRE

EXPLANATION

White Mountain plutonic-volcanic series

New Hampshire plutonic series

Oliverian plutonic series

Highlandcroft plutonic series

Ultramafic rocks

Migeosynclinal rocks

Eugeosynclinal rocks

Basement

Transition: eugeosynclinal zone to migeosynclinal zone

Unit boundary

Thrust fault — *Exact location uncertain in places, shown by stratigraphic and paleontologic evidence; sawteeth on upper plate.*

Thrust fault — *Location indicated by structural criteria; sawteeth on upper plate.*

Reverse fault — *Bars on upthrown side.*

Normal fault — *Bar and ball on downthrown side.*

Anticline

Overturned anticline

Syncline

Overturned syncline

Folds
Showing trace of axial surface, direction of plunge and approximate bearing of axis, and direction of dip of limbs; includes folds of mixed styles and three or possibly more regimes, some of which are superimposed on others at various angles. Traces are dashed where structural and stratigraphic evidence for folds is inconclusive, and are dotted where concealed by younger rocks. Anticlinoria and synclinoria are shown by lines of alternating dots and dashes where extrapolated across plutons. Anticlinoria and synforms on the flanks of domes and arches are shown by the same symbols as used for anticlines and synclines, respectively.

Trend of structural elements of basement — *Chiefly traces of axial surfaces of tight folds.*

Axial trace of recumbent fold or nappe — *Bars point into inverted limb.*

FORELAND | GREEN MOUNTAIN ANTICLINORIUM | CONNECTICUT VALLEY – GASPE SYNCLINORIUM | BRONSON HILL – BOUNDARY MOUNTAIN ANTICLINORIUM

MASSACHUSETTS

Scale of miles 0 10 20 30

TECTONIC MAP of VERMONT

CONNECTICUT VALLEY – GASPE SYNCLINORIUM | BRONSON HILL – BOUNDARY MOUNTAIN ANTICLINORIUM

TOWNSHEND–BROWNINGTON SYNCLINE | STRAFFORD–WILLOUGHBY ARCH | AVERILL SYNCLINE | GARDNER MOUNTAIN ANTICLINE

VICTORY PLUTON | LANCASTER PLUTON

West Burke | Passumpsic River | Mitchell Mt. | Burnside Mt. | Guildhall | Connecticut River

A — A′ Km. 1 Sea Level −1 −2 −3 −4 −5 −6

FORELAND | GREEN MOUNTAIN ANTICLINORIUM | CONNECTICUT VALLEY – GASPE SYNCLINORIUM | BRONSON HILL – BOUNDARY MOUNTAIN ANTICLINORIUM

CHAMPLAIN THRUST | HINESBURG SYNCLINORIUM | AXIAL ANTICLINE | TOWNSHEND–BROWNINGTON SYNCLINE | AVERILL SYNCLINE

B — Lake Champlain | Mechanicsville | HINESBURG THRUST | Huntington | Camels Hump | Green Mountains | Mud River | Winooski River | North Branch | Winooski River East Montpelier | Spruce Mountain | KNOX MOUNTAIN PLUTON | Wells River | Connecticut River | B′

Feet 5,000 Sea Level −5,000 −10,000 −15,000 −20,000

BEEKMANTOWN GROUP

Km. 1 Sea Level −1 −2 −3 −4 −5 −6

SECTION ALONG LINE B—B′

FORELAND | GREEN MOUNTAIN ANTICLINORIUM | CONNECTICUT VALLEY – GASPE SYNCLINORIUM | BRONSON HILL – BOUNDARY MOUNTAIN ANTICLINORIUM

TACONIC KLIPPEN | Taconic Range | Poultney River | Clarendon River | PINE HILL THRUST | Otter Creek | Green Mountains | Shrewsbury Pk. | Woodward Reservoir | TOWNSHEND–BROWNINGTON SYNCLINE | South Woodstock | BRATTLEBORO–AVERILL SYNCLINE | Connecticut River

C — C′ Km. 1 Sea Level −1 −2 −3 −4 −5 −6

D — D′ Feet 5,000 Sea Level −5,000 −10,000 −15,000 −20,000

MIDDLEBURY SYNCLINORIUM | AXIAL ANTICLINE | CHESTER DOME

SECTION ALONG LINE D—D′

FORELAND | G