

TABLE 1

	State of Vermont	Lamoille County
# of wells	92315	5448
# of located wells	10807	940
Mean yield, GPM	13.76	12.11
Median yield	6	6
Maximum reported yield	1200	725
Standard Deviation	22.82	19.84
Mean depth, FT	293.02	262.76
Median depth, FT	260	224
Maximum reported depth	1765	1222
Standard deviation	157.99	139.08
% wells with yield ≤ mean	70%	3919/5448 or 72%
% wells with yield > mean	30%	1529/5448 or 28%
% wells with depth ≤ mean	56%	3206/5448 or 59%
% wells with depth > mean	44%	2242/5448 or 41%

GROUNDWATER RESOURCES BY COUNTY

This county map is part of a map series used to evaluate Vermont's groundwater resources using existing data. The Lamoille County maps show yield (gallons per minute) data for bedrock wells as reported in the VT DEC Water Supply Division database. A total of 92,315 wells in the State of Vermont were analyzed in the accompanying statewide study. Data were divided into counties for presentation (Figs. 1, 2). Well locations in the database are from well driller descriptions and sketches. Some wells have been located by GPS or by correlating a well log to an E911 address. In Lamoille County, 940 out of 5,448 wells or 17% have an E911 or GPS address (Figure 3). The majority of wells, as shown on Figure 1, have suspect locations although errors due to incorrect well locations are less significant at a scale of 1:250,000.


Well yield (gpm) is generally estimated in the field with a bucket and timer. The time period is usually short and measurements are not meant to be precise. Comparisons of the mean and median values for all wells and the mean and median values for wells in Lamoille County are shown in Table 1.

Wells are grouped into yield categories on the map presented here. Depth and yield vary due to many factors, including non-geologic factors. For example, a homeowner may drill until the desired yield is obtained. The factors are not indicative of capacity. Moore et al., 2002, published "Factors Related to Well Yield in the Fractured-Bedrock Aquifer of the New Hampshire" in which they discussed a number of factors correlated positively or negatively to well yield. Among these factors are year drilled, median household income, drilling method, up gradient drainage area, thickness of overburden, depth drilled, proximity to streams/water bodies, type of bedrock, steepness of slope, elevation, fractures, and geologic structures.

The map presented is designed to be used in conjunction with other data and analyses. Groundwater flow in the crystalline bedrock of Vermont is mainly along planar features such as fractures, cleavage, faults, and bedding. These planar features may be interconnected and groundwater flow within this system is complex.

Area - specific groundwater resource studies*, available on the VGS web site, are listed below. Web: <http://www.anr.state.vt.us/dec/geo/grndwaterinx.htm>

*1. Kim, J., Springston, G., Gale, M., Dunn, R. and Becker, L., 2006. Geologic Framework for Evaluating Ground Water Resources in the Southern Worcester Mountains, Vermont. <http://www.anr.state.vt.us/dec/geo/grndwaterWorc.htm>
 2. Gale, M.H., Kim, J., King, S., Montane, P. and Orsi, C., 2006. Bedrock Geologic Map of the Southern Worcester Mountains, Vermont. Vermont Geological Survey Open File Report 2006-2.
 3. Springston, G. and Dunn, R., 2006. Surficial Geologic Map of the Southern Worcester Mountains, Vermont. Vermont Geological Survey Open File Report 2006-3.
 4. Moore, R.B., Schwarz, G.E., Clark, S.F., Jr., Walsh, G.J., and Degnan, J.R., 2002. Factors related to well yield in the fractured-bedrock aquifer of New Hampshire: USGS Professional Paper 1660.



Reported Well Yields in Bedrock Wells, Lamoille County, Vermont

by
 Marjorie Gale, George Springston, Ryan Knox and Laurence Becker
 2010

VERMONT GEOLOGICAL SURVEY

Published by:
 Vermont Geological Survey, VT DEC
 103 South Main St., Logue Cottage, Waterbury, VT 05671-2420
 Laurence Becker, State Geologist