

# **Lake Champlain Long-Term Water Quality and Biological Monitoring Program**

**Summary of Program Activities During 2017**

**March 26, 2018**

**Prepared by:**

**Vermont Department of Environmental Conservation  
Watershed Management Division  
1 National Life Drive, Main Bldg, Second Floor  
Montpelier, VT 05620-3522**

**New York State Department of Environmental Conservation  
Region 5  
P.O. Box 296  
Ray Brook, NY 12977-0296**

**Lake Champlain Research Institute  
SUNY Plattsburgh-Hudson 136D  
101 Broad Street  
Plattsburgh, NY 12901**

**Submitted to:**

**Lake Champlain Basin Program  
P.O. Box 204  
54 West Shore Rd.  
Grand Isle, VT 05458**

## **Purpose of Report**

The workplan for the Lake Champlain Long-Term Water Quality and Biological Monitoring Program approved by the Lake Champlain Basin Program specifies the following annual reporting requirements:

*An annual report will consist of a summary of the history and purpose of the (program), description of the sampling network, summary of field sampling and analytical methods, parameter listings, and data tables. The purposes of this annual report will be achieved by maintaining an up-to-date Program Description document, graphical presentations of the data, and an interactive database, including statistical summaries, on the project website..... In addition, the quarterly report produced in April each year will provide a summary of program accomplishments for the calendar year just ended, including the number of samples obtained and analyzed at each site by parameter.*

The Program Description document, interactive access to the project data, and graphical and statistical summaries of the data are available on the [program webpage](#). The purpose of this report is to provide a summary of sampling activities and other accomplishments during 2017.

## **Sampling Activities During 2017**

Table 1 lists the number of sampling visits to each lake and tributary station in relation to the target frequencies specified in the project work plan. Table 2 lists the number of samples collected and analyzed for each monitoring parameter. The New York lake and tributary field sampling was conducted by the Lake Champlain Research Institute at SUNY Plattsburgh under an MOU between NYSDEC and SUNY.

The frequency of lake sampling exceeded workplan targets at all stations during 2017. The frequency of tributary sampling was above the workplan targets for most of the stations. The number of tributary samples obtained each year depends to some extent on the number and timing of high flow events, since sampling is geared toward capturing the highest flow conditions when loading of phosphorus and other materials is greatest. Figure 1 shows that sampling at each tributary captured most peak flow events during 2017.

Table 1. Number of sampling visits during 2017 at each lake and tributary station in comparison with workplan targets.

Number of Lake Sampling Visits					Number of Tributary Sampling Visits				
Lake Station	NY	VT	Total	Workplan Target <sup>1</sup>	Tributary Station	Crew	All Parameters TP, DP, TSS, CI, TN	Total Phosphorus	Workplan Target <sup>2</sup>
2	10	9	19	12	AUSA01	NY	14	15	14/17
4	10	9	19	12	BOUQ01	NY	13	15	14/17
7	10	9	19	12	GCHA01	NY	14	15	14/17
9	10	7	17	12	LAMO01	VT	19	24	14/17
16	10	10	20	12	LAPL01	VT	14	19	14/17
19	10	10	20	12	LAUS01	NY	14	16	14/17
21	10	10	20	12	LCHA01 <sup>3</sup>	NY	14	15	14/17
25	10	10	20	12	LEWI01	VT	15	20	14/17
33	10	10	20	12	LOTT01	VT	14	18	14/17
34	10	10	20	12	METT01	VT	16	18	14/17
36	10	10	20	12	MISS01	VT	18	23	14/17
40	10	10	20	12	OTTE01	VT	16	20	14/17
46	10	9	19	12	PIKE01	VT	10	13	14/17
50	10	10	20	12	POUL01	VT	17	18	14/17
51	10	10	20	12	PUTN01 <sup>4</sup>	VT	0	0	14/17
					ROCK02	VT	14	19	14/17
					SALM01	NY	14	15	14/17
					SARA01	NY	15	16	14/17
					WINO01	VT	25	30	14/17
					JEWE02	VT	15	20	14/17
					STEV01 <sup>5</sup>	VT	13	18	14/17
					MILL01	VT	15	20	14/17

<sup>1</sup> Workplan target for lake sampling (12) applies to most chemical parameters and to phytoplankton, zooplankton, and zebra mussel veligers. Sampling for zebra mussel juveniles in Lake Champlain and for veligers in tributaries and inland lakes is done once annually.

<sup>2</sup> The project workplan calls for 14 samples per year for most chemical parameters, including 10 samples at high flow and four samples at low flow. Additional sampling for total phosphorus only should occur on 3 other dates under high flow conditions, for a target of 17 samples per year for total phosphorus.

<sup>3</sup> Little Chazy flow gage was discontinued in 2014, but was re-established on 9-25-2015.


<sup>4</sup> Putnam Creek sampling was discontinued in 2015 due to lack of funding for the flow gage.











<sup>5</sup> The USGS gage at Stevens Brook was discontinued at the end of June, 2017. A new gage was constructed by Stone Environmental and will be available online soon.


Table 2. Number of samples collected and analyzed for each monitoring parameter during 2017.

<b>Parameter</b>	<b>Lake</b>	<b>Tributaries</b>	<b>Total</b>
TP	428	450	878
DP	427	395	822
Cl	426	395	821
TN	428	451	879
Ca	59	60	119
SiO <sub>2</sub>	428	-	428
K	59	60	119
Na	59	60	119
Mg	59	60	119
Alkalinity	59	60	119
DO (Winkler)	43	-	43
Chl-a	324	-	324
TSS	-	377	377
Temperature	-	382	382
Conductivity	-	346	346
pH	-	346	346
Secchi depth	315	-	315
Multiprobe depth profiles	293	-	293
Zebra mussel veligers	121	-	121
Zebra mussel settled juveniles	4	-	4
Mysids	94		94
Zooplankton	150		150
Phytoplankton	135		135
Spiny waterflea	454		454

Figure 1. Sampling dates during 2017 in relation to daily flows at each tributary station. Daily flows are shown by lines, and sampling dates are shown by dots.







## Data Quality Assurance Results

As described in the program's Quality Assurance Project Plan, field equipment blanks and field duplicate samples are obtained on each sampling run. The results for the blank samples are summarized in Table 3. Thirteen of the 277 blank samples analyzed during 2017 (4.7%) had concentrations above the analytical detection limits. Results for field duplicate samples are summarized in Table 4 for the chemical analyses.

The results from laboratory and field duplicate analyses run on phytoplankton samples obtained during 2011-2015 are shown in Table 5. Phytoplankton data are not yet finalized for 2017.

Table 3. Field equipment blank results during 2017 for lake and tributary samples.

Test	Detection Limit	Units	Number of Blanks Obtained	Number of Blanks Above Limit	High Blank Values
Alk	1.0	mg/l	6	0	
Cl	2.0	mg/l	42	0	
TN	0.1	mg/l	45	0	
TP	5.0	µg/l	45	2	6.3 47
DP	5.0	µg/l	42	6	5.1 5.5 7.1 11.4 13.5 45.3
Chl-a	0.5	µg/l	14	3	.94 3.7 8.8
TSS	1.0	mg/l	26	1	33.5
SiO <sub>2</sub>	0.2	mg/l	15	0	
Al	20	µg/l	6	0	
Fe	50	µg/l	6	0	
Ca	0.1	mg/l	6	1	.105
Na	0.5	mg/l	6	0	
K	0.5	mg/l	6	0	
Mg	0.02	mg/l	6	0	
Mn	5	µg/l	6	0	
Total			277	13	

Table 4. Field duplicate results for chemical tests during 2017 showing the number of duplicates obtained (N) and the mean relative percent difference (RPD) between duplicate pairs.

<b>Test</b>	<b>N</b>	<b>Mean RPD</b>
Chl-a	16	25.2
Cond	--	--
Cl	48	2.1
DP	48	11.4
pH	--	--
Alk	6	.8
TN	49	5.8
TP	50	9.2
TSS	31	9.4
SiO <sub>2</sub>	16	1.4
Al	6	10.8
Ca	6	.9
Fe	6	7.6
K	6	1.1
Na	6	1.2
Mg	6	.9
Mn	6	6.1

## **Phytoplankton and Zooplankton Database**

All phytoplankton data from 2006-2015 have been incorporated into the main Lake Champlain Monitoring Program database. Zooplankton data are currently available for the project period of 1993-2013. The data available for download from the web interface include phytoplankton cell densities and biovolumes, and zooplankton densities grouped by major taxonomic category. Counts by individual taxa will eventually be added to the web page, but are currently available only by request.



Table 5. Phytoplankton duplicate results for 2011–2015, the most recent sampling data, showing the number of pairs (N) and the mean relative percent difference (RPD) between pairs.

Test	Year	N	Sample Type	Mean RPD
Field Duplication	2011	11	Biovolume	44.4
			Cell Density	44.8
	2012	12	Biovolume	42.3
			Cell Density	55.4
	2013	8	Biovolume	27.9
			Cell Density	33.4
	2014	8	Biovolume	37.8
			Density	35.3
	2015	10	Biovolume	53.0
			Density	71.5
Lab Duplication	2011	15	Biovolume	32.1
			Cell Density	39.3
	2012	19	Biovolume	28.3
			Cell Density	27.5
	2013	12	Biovolume	26.9
			Cell Density	37.1
	2014	13	Biovolume	48.2
			Cell Density	51.7
	2015	14	Biovolume	32.6
			Density	30.3

## Wastewater Phosphorus Discharge Data

The project workplan requires an annual compilation of wastewater phosphorus discharge data for all treatment facilities in the Vermont and New York portions of the Lake Champlain Basin. Data on annual mean flow, total phosphorus concentration, and phosphorus load at each facility have been compiled for 2016 along with data from previous years, and are available electronically in spreadsheet form on request. The total loads and flows from Vermont and New York wastewater treatment facilities during 2007-2016 are summarized in Table 6. Vermont 2017 data is delayed due to new online reporting requirements.

Table 6. Total phosphorus load to Lake Champlain from wastewater treatment facilities in Vermont and New York from 2007-2016.

State	Number of Facilities	Year	Total Phosphorus Load (mt/yr)	Total Flow (mgd)
Vermont	60	2007	20.9	43.5
	60	2008	21.1	45.1
	60	2009	20.3	40.5
	60	2010	18.4	39.7
	59	2011	19.3	45.5
	59	2012	16.9	37.6
	59	2013	17.1	40.6
	59	2014	18.8	40.7
	59	2015	13.6	38.5
	59	2016	11.7	36.5
New York	29	2007	28.5	33.2
	29	2008	26.5	34.3
	29	2009	20.9	31.5
	29	2010	22.0	32.8
	29	2011	23.0	34.4
	29	2012	22.6	30.4
	29	2013	22.9	30.3
	29	2014	24.7	30.3
	29	2015	23.7	29.6
	29	2016	22.2	30.2

### Rock River Monitoring Project

A Rock River Watershed Targeted Best Management Practice (BMP) Implementation Project was initiated in 2010 with funding provided by the Lake Champlain Basin Program (LCBP) and with oversight provided by a coordinating committee including the U.S. Natural Resource Conservation Service (NRCS), the Vermont Agency of Agriculture, Food, and Markets (AAFV) and the Vermont Department of Environmental Conservation (DEC). The purpose of the project is to demonstrate water quality improvements from a focused agricultural BMP implementation effort in a small watershed where very high rates of phosphorus loading to Lake Champlain have been documented. BMP implementation got underway in 2010 and new installations are added each year.

In order to document water quality improvements resulting from the targeted BMP implementation in the Rock River watershed, the Vermont DEC established monitoring stations immediately upstream and downstream of the BMP implementation area in late 2010 and funded the construction and operation of a U.S. Geological Survey (USGS) stream flow gage at the downstream site. The DEC issues grants to the Friends of Northern Lake Champlain (FNLC) to support sample collection activities by trained local residents, and the DEC Laboratory conducts the sample analyses. The LCBP financially supports the laboratory analytical efforts and supported the stream gaging through September 2014. The State of Vermont now supports the Rock River stream gage through a cooperative agreement with the USGS.

The area targeted for BMP implementation is approximately 13.5 km<sup>2</sup> in size on the upper Rock River in the towns of Highgate and Franklin, VT. Nearly all of the BMP implementation area is contained within the catchment area between the upstream monitoring station (RR20) and the downstream station (RR14). The implementation area occupies about 90% of the 15.1 km<sup>2</sup> drainage area between the upstream and the downstream stations. A USGS continuous stream flow gage is co-located with the downstream sampling station (RR14).

Samples are obtained manually as grab samples from the center of the river on each date for analysis of total phosphorus (TP), total dissolved phosphorus (DP), and total suspended solids (TSS). Sampling is conducted biweekly year-round except during the winter months when snow and ice in the river make sampling impossible. Additional sampling is conducted during high-flow events.


The study was designed as an upstream/downstream, before/after analysis, which is a type of a paired watershed design (Clausen and Spooner, 1993). It was originally anticipated that Analysis of Covariance (ANCOVA) would be used to test for statistically significant differences in TP, DP, PP, and TSS concentrations at the downstream station before vs. after BMP implementation (Meals, 2004). However, agricultural BMP implementation did not occur during a discrete, short-term time interval that would allow for a clear distinction between pre and post-implementation periods. Consequently, regression analysis of continuous temporal trends was used in place of the original ANCOVA design.

There have been 359 upstream/downstream paired samples collected and analyzed for TP, DP, and TSS through 2017. This total includes some samples obtained during 2008-2009 by Vermont DEC as part of a previous study. The numbers of paired samples obtained each year are shown in Table 7. The project site map is shown in Figure 2.

Table 7. Numbers of paired samples obtained.

Year	Number of sample pairs
2008	10
2009	2
2010	18
2011	66
2012	55
2013	55
2014	51
2015	27
2016	33
2017	42
<b>Total</b>	<b>359</b>

Figure 2– Rock River Monitoring Project site map



Late in 2017, through discussions about National Water Quality Initiative (NWQI) activities in the watershed, we learned that BMP projects were also installed in the control watershed beginning in 2013 when the Rock River became a priority for the NRCS (Figure 3). As a result, the original study design has been rendered invalid. However, a steering committee including representation from VT DEC, NRCS, VAAFM, LCBP, and MDDELCC formed in early 2018 felt strongly that there was value that could be obtained from continued monitoring of the downstream station (#14). This station is influenced by runoff from the original control and treatment watersheds and can be used to monitor water quality changes over time in response to existing and continued BMP implementation. This information will still be useful to understand the success of management efforts. During 2018, monitoring will continue at station 14 and the steering committee will update the study design to include all BMP activities upstream of this site.




Figure 3. Location of BMP implementation sites funded by NWQI in the two subwatersheds targeted by this project. (Image provided by the NRCS.)

### Invasive Species Monitoring Lake Champlain

A total of 454 zooplankton samples were scanned for *Bythotrephes longimanus* (spiny waterflea) from monitoring stations on Lake Champlain in 2017 (Table 8, Figures 4.1 – 4.4). Whole water vertical tows were taken at each monitoring station using a 250 µm mesh 50 cm plankton net. In addition to the whole water tows, an epilimnion tow was collected at sites 7, 9, 19, 25, 34 and 36 for each sampling event. If the sites were isothermic, epilimnion tows were taken at 2x Secchi depth or 1m from the bottom. Samples were then taken to the laboratory where they were visually scanned under a dissecting microscope to determine population densities. All samples were also scanned for other potential invasive invertebrates including *Hemimysis anomala* and *Cercopagis pengoi*. Based on whole water vertical tows, densities of *B. longimanus* notably decreased in 2017, with only two individuals recorded in May (Figure 4.1) and one in

Table 8. Spiny water flea (SWF) monitoring stations in the Lake Champlain.				# of	
62			4	24	
51	45.0410	73.1290	10	20	
50	45.0130	73.1740	10	20	
46	44.9480	73.3400	10	20	
40	44.7850	73.1620	10	20	
36	44.7560	73.3350	10	30	
34	44.7080	73.2270	10	30	
33	44.7010	73.4180	10	20	
25	44.5820	73.2810	10	30	
21	44.4740	73.2320	10	20	
19	44.4710	73.2990	10	78	
16	44.4250	73.2220	10	20	
10	44.3000	73.3214	4	22	
9	44.2420	73.3340	10	30	
7	44.1260	73.4120	10	31	
4	43.9540	73.4050	10	20	
2	43.7140	73.3830	10	20	
			Total # of Samples	454	

September (Figure 4.3) for a total of three specimens in 2017.




Figure 4.1 Spiny water flea density from vertical whole water and epilimnion tows from May-June 2017




Figure 4.2 Spiny water flea density from vertical whole water and epilimnion tows from July- Early August 2017




Figure 4.3 Spiny water flea density from vertical whole water and epilimnion tows from Late August– Late September 2017





Figure 4.4 Spiny water flea density from vertical whole water and epilimnion tows from October 2017



Figure-5 Lake Champlain  
LTM Sampling Locations

